

Presentation

Each film coated Neurex-B $^{\otimes}$ tablet contains Thiamine Mononitrate (B₁) USP 100 mg, Pyridoxine Hydrochloride (B₆) USP 200 mg, Cyanocobalamin (B₁₂) USP 200 mcg.

Indications

Neurex-B[®] is administered as a part of combine therapy in B_1 , B_6 , B_{12} deficiency diseases. The drug is also indicated in the treatment of Chronic pain, Diabetic neuropathy, Peripheral neuralgia, Lumbago, Myalgia, Sciatica, Facial neuralgia, Intercostal neuralgia & Spinal pain.

Dosage and Administration

Daily dose 1 to 3 tablets as directed by the physician.

Side Effects

Very rare allergic reactions, in high doses nervous excitement may be seen.

Cautions

Cyanocobalamin should not be given before a diagnosis has been fully established because of the possibility of masking symptoms of subacute degeneration of the spinal cord.

Contraindications

Drug should not be used in the patients with Levodopa therapy and hypersensitivity to any of the active ingredients.

Drug Interactions

Pyridoxine decreases the antiparkinsonic effect of Levodopa.

Use in Pregnancy and Breast-Feeding

Neurex-B $^{\! ^{\otimes_{\! -}}}$ is not contraindicated but tablet can be given as directed by the physician.

Overdose

Nervous excitement, tachycardia, pain in the heart may occur due to overdose. So treatment should be symptomatic and supportive.

Commercial Pack

Each box contains 3 blister strips of 10 tablets.

