

# Sputen<sup>®</sup>

**Bromhexine HCl**

## Presentation

Sputen<sup>®</sup> Tablet : Each tablet contains Bromhexine HCl BP 8 mg  
Sputen<sup>®</sup> Oral Solution : Each 5ml oral solution contains Bromhexine HCl BP 4mg

## Indications

Sputen<sup>®</sup> is indicated in the treatment of respiratory disorders associated with productive cough. These include; tracheobronchitis, bronchitis with emphysema, bronchiectasis, bronchitis with bronchospasm, chronic inflammatory pulmonary conditions and pneumoconiosis.

## Dosage and Administration

### Sputen<sup>®</sup> Tablet

Adults and children over 10 years : 8-16 mg 3 times daily.

Children 5-10 years:

### Sputen<sup>®</sup> oral solution

Adults : The recommended daily dose is 2 to 4 teaspoonful 3 times Initially 4 teaspoonful 3 times daily and then as required.

Children : Suggested dosage for children under 2 years is ¼ teaspoonful 3 times daily, for 2-5 years ½ teaspoonful 3 times daily and for children aged 5-10 years 1 teaspoonful 3 times Daily.

## Side effects

Gastrointestinal side effects may occur occasionally with bromhexine and a transient rise in serum aminotransferase values has been reported. Other reported adverse effects include headache, dizziness, sweating and skin rash.

## Precautions

Since mucolytics may disrupt the gastric mucosa so bromhexine should be used with care in patients with a history of peptic ulceration. Care is also advisable in asthmatic patients.

## Contraindication

Contraindicated to those who are hypersensitive to bromhexine hydrochloride.

## Storage instructions

Store below 25°C. Protect from light. Keep the container tightly closed.

## Commercial Pack

Sputen<sup>®</sup> Tablet: Each Box contains 10 blister Pack 10 tablets..

Sputen<sup>®</sup> Oral Solution : Bottle Contains 100ml oral solution.

Manufactured by :  
 **Silva**  
**Pharmaceuticals Limited**  
Majdee, Noakhali, Bangladesh